

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.


JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Another interesting expedition was that to *Tiahuanuco* and the sacred islands of Incas, on lake *Titicaca*, which led to results of much archæological interest. Important aid has been rendered to the expedition by many residents in Peru without whose assistance the establishment of the station would have been extremely difficult.

HARVARD COLLEGE OBSERVATORY, CAMBRIDGE, MASS., February 19, 1892.


### ASTRONOMICAL OBSERVATIONS.


Made by Torvald Köhl, at Odder, Denmark, in the year 1891.



The region near the star 26 Cygni has been reviewed. December 7, 1890, the star Z appeared as bright as a, but it very rapidly decreased, so that on

```
January
           I: Z < c.
           6: id.
  ..
           9: id.
February 4: Z invisible in the 3-inch Steinheil, power 42.
          13: id.
  "
          27: id.
          26: Z = b.
August
          30: id.
September 4: a > Z < c.
           8: Z = c.
          23: Z < c.
          29: Z = e.
October
         26: Z invisible.
November 5: Z < e, extremely faint.
          28: id.
```





This variable has been observed at the following times. On December 7, 1890, the star was invisible, but it must have rapidly grown brighter, the maximum being calculated to fall on February 1, 1891.

```
February
           4: T a little fainter than a.
          13: a > T > c; T perhaps a little brighter than b.
          17: id.
          26: id., b almost = c.
March
           9: id.
April
           3: T a little fainter than d.
September 10: T almost invisible, fainter than g.
          23: e < T < d.
          24: id.
          29: T = d.
November 6: T > a.
                        N. B.
          28: T = b.
```

In "Publications A. S. P.," Vol. III, No. 15, page 101, I directed attention to a little star (15) near 61 Cygni, which twenty years ago was seen by Professor Rudolph Falb, and at the present time is not brighter than 13½ magnitude. I have often looked out for this little star thinking it might reappear as a star of the 11th magnitude, but without success. Many other parts of constellations, where variable stars were supposed to be situated, have also been reviewed, the region around Tycho's Stella nova Cassiopeiæ, for example.

## The Solar Eclipse of June 6.

This partial eclipse, which, on account of the bad weather could not be observed at the Observatory of Copenhagen, was seen here in the following phases:

At 5 38 P. M. (time of Copenhagen), the eclipse begins about in the middle of the N. W. quadrant.

- 6 o "The disc of the moon is crossing the sun's disc in the north point of the latter.
- 6 11 15 "The limb of the moon is touching a large sun-spot in the position

$$\triangle a = +0.8'$$

$$\triangle \delta = +6.0'$$


- 6 30 "Maximum of the eclipse (0.4).
- 6 47 30 " Emersion of the above named sun-spot.
- 7 o "At this moment only 0.14 of the sun's diameter is darkened.
- 7 14 "End of the eclipse.

At the middle of the observation the sun's altitude was about 14°, and the limit of the moon therefore was very undulating.

## The Lunar Eclipse of November 15.

h. I I	m. 25	s.	Р. М.	The shadow is touching the eastern limit of the
				moon in 22° northern latitude.
11	32		"	• The shadow crossing Aristarchus.
ΙΙ	36		"	The shadow touches the southern wall of Grimaldi.
11	44		"	The shadow through Copernicus.
ΙI	45		"	The shadow through Plato.
II	47		"	The shadow through Archimedes.
11	54		"	The shadow has reached the eastern wall of
				Aristoteles and Eudoxus.
ΙI	59		"	The shadow crossing Bessel, Menelaus and
				Thebit.
I 2	4		"	The shadow is passing <i>Posidonius</i> , <i>Plinius</i> and
				the eastern wall of Tycho.
12	5	30	"	The shadow is passing the western wall of <i>Tycho</i> .
I 2	II		"	The shadow through <i>Proclus</i> and the western wall of <i>Clavius</i> and <i>Theophilus</i> .
12	27		"	Totality of the eclipse.

No further observations possible, as heavy clouds were crossing the sky.



## Sun-Spot Observations,

On the following days the sun has been observed:

January 6, 14, 18.

February 9, 12, 13, 15, 16, 17, 18, 19, 24, 26, 27, 28.

March 2, 7, 17, 18, 19, 21, 22, 26, 31.

April 3, 4, 5, 6, 9, 10, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29.

May 2, 3, 4, 5, 6, 8, 10, 11, 12, 13, 14, 15, 18, 19, 23, 24, 25, 27, 28, 29, 30, 31.

June 1, 2, 5, 6, 7, 8, 9, 10, 11, 12, 15, 16, 20, 21, 22, 29, 30. July 1, 6, 8, 10, 12, 13, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31.

August 1, 3, 5, 6, 7, 8, 9, 10, 12, 15, 16, 17, 19, 21, 22, 24, 26, 28, 29, 30, 31.

September 2, 3, 7, 9, 11, 13, 14, 19, 23, 24, 25, 27, 29, 30. . October 1, 3.

A number of (44) sketches has been made and positions of a great many sun-spots obtained. In fig. 3, the numbers at the edge indicate differences of R. A. and Decl. in parts; the sun's diameter being divided into 40 parts.

#### Meteors.

No.	Time.	Beginning.	End.	Mag- nitude.	Notes.
I	h. m. s. Feb. 4, 8 6 0	167 + 34	182 + 36	2	Slow.
2	Aug. 11, 10 20 0	255 + 42	272 + 27	I	
3	" 34 0	313 + 30	290 + 15	I	
4	". 38 30	262 + 15	262 + 5	1	
5	" 44 0	315 + 34	300 + 24	2	
6	" 56 o	311 + 37	297 + 27	2	
7	" 58 15	354 + 18	347 + 7	2	
8	" II 2 0	338 + 23	323 + 10	3	Train.

Cloudy weather troubled the observations, which were planned in connection with station Copenhagen, on the 9th, 10th and 12th of August. Occasionally, while observing Wolf's comet on September 8, a bright shooting star happened to cross the field at 9<sup>h</sup> 50<sup>m</sup>. The comet was about 1° S. E. from 64 *Pleiadum*.

In the year 1891 a number of (13) fire-ball observations from different places in Denmark has been registered.